

## IIT-JEE 2010

## Chemistry Part - I

### PART I - Chemistry

#### SECTION - I

#### Single Correct Choice Type

This Section contains **8 multiple choice questions**. Each question has four choices (A), (B), (C) and (D) out of which **ONLY ONE** is correct.

1. The bond energy (in kcal mol<sup>-1</sup>) of a C–C single bond is approximately  
 (A) 1                      (B) 10                      (C) 100                      (D) 1000

1. **(C)** It is a fact. It is nearly 84 kcal mol<sup>-1</sup>

2. The species which by definition has ZERO standard molar enthalpy of formation at 298 K is

(A) Br<sub>2</sub>(g)                      (B) Cl<sub>2</sub>(g)                      (C) H<sub>2</sub>O(g)                      (D) CH<sub>4</sub>(g)





2. **(B)** Cl<sub>2</sub>(g) is gas in elemental state Br<sub>2</sub> is not of because it is liquid in elemental state.

3. The ionization isomer of [Cr(H<sub>2</sub>O)<sub>4</sub>Cl(NO<sub>2</sub>)]Cl is

(A) [Cr(H<sub>2</sub>O)<sub>4</sub>(O<sub>2</sub>N)]Cl<sub>2</sub>                      (B) [Cr(H<sub>2</sub>O)<sub>4</sub>Cl<sub>2</sub>](NO<sub>2</sub>)  
 (C) [Cr(H<sub>2</sub>O)<sub>4</sub>Cl(ONO)]Cl                      (D) [Cr(H<sub>2</sub>O)<sub>4</sub>Cl<sub>2</sub>(NO<sub>2</sub>)]H<sub>2</sub>O






3. **(B)** [Cr(H<sub>2</sub>O)<sub>4</sub>Cl(NO<sub>2</sub>)]Cl and [Cr(H<sub>2</sub>O)<sub>4</sub>Cl<sub>2</sub>](NO<sub>2</sub>) are ionization isomers.


4. The Correct structure of ethylenediamineteraacetic acid (EDTA) is



4. **(C)** It is a fact






8. In the reaction   $\xrightarrow{\text{HBr}}$  the products are
- (A)  and  $\text{H}_2$  (B)  and  $\text{CH}_3\text{Br}$
- (C)  and  $\text{CH}_3\text{OH}$  (D)  and  $\text{CH}_3\text{Br}$



**SECTION - II**  
**Multiple Correct Choice Type**



This section contains 5 multiple choice questions. Each question has four choices (A), (B), (C) and (D) out of which ONE OR MORE may be correct.

9. In the reaction   $\xrightarrow{\text{NaOH(aq)} / \text{Br}_2}$  the intermediate(s) is (are)




10. The reagent(s) used for softening the temporary hardness of water is (are)
- (A)  $\text{Ca}_3(\text{PO}_4)_2$  (B)  $\text{Ca}(\text{OH})_2$  (C)  $\text{Na}_2\text{CO}_3$  (D)  $\text{NaOCl}$
10. (BC)  $\text{Na}_2\text{CO}_3 + \text{Ca}(\text{HCO}_3)_2 \rightarrow \text{CaCO}_3 + \text{NaHCO}_3$   
 $\text{Ca}(\text{OH})_2 \rightarrow \text{CaO} + \text{H}_2\text{O}$   
 $\text{CaO} + \text{Ca}(\text{HCO}_3)_2 \rightarrow 2\text{CaCO}_3 + \text{H}_2\text{O}$

11. Aqueous solutions of  $HNO_3$ ,  $KOH$ ,  $CH_3COOH$ , and  $CH_3COONa$  of identical concentrations are provided. The pair(s) of solutions which form a buffer upon mixing is (are)
- (A)  $HNO_3$  and  $CH_3COOH$  (B)  $KOH$  and  $CH_3COONa$   
 (C)  $HNO_3$  and  $CH_3COONa$  (D)  $CH_3COOH$  and  $CH_3COONa$




12. In the Newman projection for 2,2-dimethylbutane



X and Y can respectively be

- (A) H and H (B) H and  $C_2H_5$  (C)  $C_2H_5$  and H (D)  $CH_3$  and  $CH_3$




13. Among the following, the intensive property is (properties are)
- (A) Molar conductivity (B) electromotive force  
 (C) resistance (D) heat capacity
13. (AB) resistance and heat capacity both depend on mass.

### SECTION - III Paragraph Type

This section contains 2 paragraphs. Based upon the first paragraph 2 multiple choice questions and based upon the second paragraph 3 multiple choice questions have to be answered. Each of these questions has four choices (A), (B), (C) and (D) out of which ONLY ONE is correct.

#### Paragraph for questions 14 to 15

The concentration of potassium ions inside a biological cell is at least twenty times higher than the outside. The resulting potential difference across the cell is important in several processes such as transmission of nerve impulses and maintaining the ion balance. A simple model for such a concentration cell involving a metal  $M$  is :



For the above electrolytic cell the magnitude of the cell potential  $|E_{\text{cell}}| = 70 \text{ mV}$ .

14. For the above cell
- (A)  $E_{\text{cell}} < 0; \Delta G > 0$  (B)  $E_{\text{cell}} > 0; \Delta G < 0$   
 (C)  $E_{\text{cell}} < 0; \Delta G^\circ > 0$  (D)  $E_{\text{cell}} > 0; \Delta G^\circ < 0$
14. (B) The reaction should be spontaneous and hence  $\Delta G < 0$ .

15. If the 0.05 molar solution of  $M^+$  is replaced by a 0.0025 molar  $M^+$  solution, then the magnitude of the cell potential would be  
 (A) 35 mV (B) 70 mV (C) 140 mV (D) 700 mV

15. (C)  $0.070 = Emf^\circ - \frac{.06}{1} \log .05$

$$x = Emf^\circ - \frac{.06}{1} \log .0025$$

$$\frac{.070}{x} = \frac{\log .05}{\log .0025} = \frac{-1.3}{-2.6}$$

$$x = .07 \times 2 \times 10^3 = 140$$

### Paragraph for questions 16 to 18

Copper is the most noble of first row transition metals and occurs in small deposits in several countries. Ores of copper include chalcocite ( $Cu_2O$ ), malachite ( $Cu_2(OH)_2CO_3$ ), azurite ( $Cu_3(CO_3)_2(OH)_2$ ), and chalcocyanite ( $Cu_5(CO_3)_4(OH)_2$ ). However, 80% of the world copper production comes from the ore chalcopyrite ( $CuFeS_2$ ). The extraction of copper from chalcopyrite involves partial roasting, removal of iron and self-reduction.

16. Partial roasting of chalcopyrite produces  
 (A)  $Cu_2S$  and  $FeO$  (B)  $Cu_2O$  and  $FeO$   
 (C)  $CuS$  and  $Fe_2O_3$  (D)  $Cu_2O$  and  $Fe_2O_3$
16. (A) Partial roasting of  $CuFeS_2$  gives  $Cu_2S$  and  $FeO$ . Iron is preferentially oxidised.
17. Iron is removed from chalcopyrite as  
 (A)  $FeO$  (B)  $FeS$  (C)  $Fe_2O_3$  (D)  $FeSiO_3$
17. (D)  $FeO + SiO_2 \rightarrow FeSiO_3 \downarrow$
18. In self-reduction, the reducing species is  
 (A) S (B)  $O^{2-}$  (C)  $S^{2-}$  (D)  $SO_2$
18. (C)  $Cu_2S + 2Cu_2O \rightarrow 6Cu + SO_2$

### SECTION - IV (Integer Type)


This section contains TEN questions. The answer to each question is a single digit integer ranging from 0 to 9. The correct digit below the question number in the ORS is to be bubbled.

19. The number of neutrons emitted when  ${}_{92}^{235}U$  undergoes controlled nuclear fission to

${}_{54}^{142}Xe$  and  ${}_{38}^{90}Sr$  is :

19. (3)  ${}_{92}^{235}U \rightarrow {}_{54}^{142}Xe + {}_{38}^{90}Sr + {}_0^1n$   
 $235 = 232 + x$   
 $x = 3.$


20. The total number of basic groups in the following form of lysine is




20. (2)  $-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}^{\ominus}$  is Conjugate base,  $-\text{NH}_2$  is basic,  $\text{NH}_3^+$  is acidic

Total no. is 2.

21. The total number of cyclic isomers possible for a hydrocarbon with the molecular formula  $\text{C}_4\text{H}_6$  is.




22. In the scheme given below, the total number of intramolecular aldol condensation product formed from 'Y' is




22. (1) Since it is symmetrical only one aldol product is possible.

23. Amongst the following, the total number of compounds soluble in aqueous  $\text{NaOH}$  is




23. (4)  $\text{COOH}$  group and  $\text{OH}$  group reacts with  $\text{NaOH}$ .

24. Amongst the following, the total number of compounds whose aqueous solution turns red litmus paper blue is



24. (3)  $\text{KCN}$ ,  $\text{K}_2\text{CO}_3$  and  $\text{LiCN}$  They are the salts of weak acid and strong base.

25. Based on VSEPR theory, the number of 90 degree  $F-B-F$  angles in  $BrF_5$  is



Due to  $LP-BP$  repulsion the bond angle decreases axial angle no longer remains  $90^\circ$  other angle also suffers distortion.

26. The value of  $n$  in the molecular formula  $Be_nAl_2Si_6O_{18}$  is

26. (3)  $2n + 2 \times 3 + 6 \times (4) + 18(-2) = 0$   $2n = 6$   $n = 3$

27. A student performs a titration with different burettes and finds titre values of 25.2 mL, 25.25 mL, and 25.0 mL. The number of significant figures in the average titre value is

27. (3) As per rules

28. The concentration of  $R$  in the reaction  $R \rightarrow P$  was measured as a function of time and the following data is obtained :

$[R]$ (molar)	1.0	0.75	0.40	0.10
t(min.)	0.0	0.05	0.12	0.18

The order of the reaction is

28. (0)  $(.25 / .05) = 5$  again  $(0.35 / 0.07) = 5$   
rate remain constant irrespective of change of concentration.